



(11) **EP 1 498 484 A1**

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:
19.01.2005 Bulletin 2005/03

(51) Int Cl. 7: **C12N 15/09, C12N 9/04,**
C12N 1/15, C12N 1/19,
C12N 1/21, C12N 5/00

(21) Application number: **03719216.8**

(86) International application number:
PCT/JP2003/005375

(22) Date of filing: **25.04.2003**

(87) International publication number:
WO 2003/091430 (06.11.2003 Gazette 2003/45)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR
Designated Extension States:
AL LT LV MK

(72) Inventor: **Sode, Koji**
Tokyo 152-0013 (JP)

(30) Priority: **26.04.2002 JP 2002125353**

(74) Representative: **Bannerman, David Gardner et al**
Withers & Rogers,
Goldings House,
2 Hays Lane
London SE1 2HW (GB)

(71) Applicant: **Sode, Koji**
Tokyo 152-0013 (JP)

(54) **GLUCOSE DEHYDROGENASE BETA-SUBUNIT AND DNA ENCODING THE SAME**

(57) A DNA fragment encoding a β subunit is obtained by inverse PCR using primers designed based on the nucleotide sequence of a N-terminal signal sequence region of a GDH β subunit derived from *Burkholderia cepacia* KS1 strain.

EP 1 498 484 A1

Printed by Jouve 75001 PARIS (FR)

BEST AVAILABLE COPY

EP 1 498 484 A1

Description

Technical Field

- 5 [0001] The present invention relates to cytochrome C that constitutes a glucose dehydrogenase β subunit, to a DNA encoding the cytochrome C, and to their utilization. The glucose dehydrogenase is useful for a glucose sensor using an enzyme electrode or the like.

Background Art

- 10 [0002] Biosensors using an enzyme that specifically reacts with a particular substrate are being actively developed in various industrial fields. As for a glucose sensor, which is one of the biosensors, in particular, measurement methods and devices utilizing such methods are being actively developed mainly in medical fields. For example, the glucose sensor has a history of about 40 years since Clark and Lyons first reported about a biosensor including glucose oxidase and an oxygen electrode in combination in 1962 (L.C. Clark, J. and Lyons, C. "Electrode systems for continuous monitoring in cardiovascular surgery." Ann. n. y. Acad. Sci., 105: 20-45).

- 15 [0003] Thus, the adoption of glucose oxidase as an enzyme for the glucose sensor has a long history. This is because glucose oxidase shows high substrate specificity for glucose and superior thermal stability, this enzyme can further be produced in a large scale, and its production cost is lower than those of other enzymes. The high substrate specificity means that this enzyme does not react with a saccharide other than glucose, and this leads to an advantage that accurate measurement can be achieved without error in measurement values. Further, the superior thermal stability means that problems concerning denaturation of the enzyme and inactivation of its enzymatic activity due to heat can be prevented, and this leads to an advantage that accurate measurement can be performed over a long period of time.

- 20 [0004] However, although glucose oxidase has advantages as described above, it has a problem that the enzyme is affected by dissolved oxygen and this affects measurement results.

- 25 [0005] Meanwhile, in addition to glucose oxidase, a glucose sensor utilizing glucose dehydrogenase (hereinafter referred to as "glucose dehydrogenase" or "GDH") has also been developed. This enzyme is also found in microorganisms. For example, there are known glucose dehydrogenase derived from *Bacillus* (EC 1.1.1.47) and glucose dehydrogenase derived from *Cryptococcus* (EC 1.1.1.119).

- 30 [0006] The former glucose dehydrogenase (EC 1.1.1.47) is an enzyme that catalyzes a reaction of β -D-glucose + NAD(P)⁺ \rightarrow D- δ -gluconolactone + NAD(P)H + H⁺, and the latter glucose dehydrogenase (EC 1.1.1.119) is an enzyme that catalyzes a reaction of D-glucose + NADP⁺ \rightarrow D- δ -gluconolactone + NADPH + H⁺. The aforementioned glucose dehydrogenases derived from microorganisms are already marketed.

- 35 [0007] These glucose dehydrogenases have an advantage that they are not affected by dissolved oxygen in a measurement sample. This leads to an advantage that accurate measurement can be achieved without causing errors in measurement results even when the measurement is performed in an environment in which the oxygen partial pressure is low, or a high-concentration sample requiring a large amount of oxygen is used for the measurement.

- 40 [0008] However, although conventional glucose dehydrogenase is not affected by dissolved oxygen, it has problems of poor thermal stability and substrate specificity poorer than that of glucose oxidase. For an enzyme which is used in a sensor, an enzyme that overcomes disadvantages of both of glucose oxidase and glucose dehydrogenase has been desired.

- 45 [0009] The inventors of the present invention reported results of their studies about GDH using samples collected from soil near hot springs in Sode, K., Tsugawa, W., Yamazaki, T., Watanabe, M., Ogasawara, N., and Tanaka, M., Enzyme Microb. Technol., 19, 82-85 (1996); Yamazaki, T., Tsugawa, W. and Sode, K., Appl. Biochem. and Biotec., 77-79/0325 (1999); and Yamazaki, T., Tsugawa, W. and Sode, K., Biotech. Lett., 21, 199-202 (1999). The microorganisms in those samples produce a coenzyme-binding GDH, and the enzymologic properties such as optimum reaction temperature, thermal stability, and substrate specificity have already been clear (See the aforementioned documents). This enzyme is a hetero oligomeric enzyme that is constituted by a catalyst subunit having a high thermal resistance (α subunit), an electron transferring subunit (β subunit), and γ subunit having an unknown function, and the activity peaks thereof are observed at 45°C and 75°C, respectively. Further, the γ and α subunit genes have been cloned, and it has been clarified that the aforementioned microorganism belongs to *Burkholderia cepacia*, and the N-terminal amino acid sequence of the β subunit has been clarified (Ken Inose, Tokyo Agricultural Engineering University Master's Thesis (2001)). However, the structure of the β subunit gene has not been reported.

55 Disclosure of the Invention

- [0010] An object of the present invention is to provide a DNA encoding a GDH β subunit of a microorganism belonging to the genus *Burkholderia* and a method of using the DNA.

EP 1 498 484 A1

[0011] The inventors of the present invention have further advanced the study on GDH of *Burkholderia cepacia* KS1 strain and were successful in isolating a DNA encoding a GDH β subunit, thereby completing the present invention.

[0012] That is, the present invention can be described as follows.

(1) A protein defined in the following (A) or (B):

- (A) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16;
(B) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16 including substitution, deletion, insertion or addition of 1 to 20 amino acid residues.

(2) A DNA encoding a protein defined in the following (A) or (B):

- (A) a protein which has at least the amino acid sequence consisting of amino acids 23 to 425 of SEQ ID NO: 16;
(B) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16 including substitution, deletion, insertion or addition of 1 to 20 amino acid residues.

(3) The DNA according to item (2), in which the DNA is defined in the following (a) or (b):

- (a) a DNA including the nucleotide sequence consisting of nucleotides 187 to 1398 of SEQ ID NO: 15;
(b) a DNA which is hybridizable with the nucleotide sequence consisting of nucleotides 187 to 1398 of SEQ ID NO: 15 under stringent conditions.

(4) The DNA according to item (3), further including the nucleotide sequence consisting of nucleotides 121 to 187 of SEQ ID NO: 15.

(5) A recombinant vector including a DNA according to any one of items (2) to (4).

(6) A transformant transformed with a DNA according to any one of items (2) to (4) or the recombinant vector according to item (5).

(7) A method of producing a glucose dehydrogenase β subunit, including culturing the transformant according to item (6) to produce a glucose dehydrogenase β subunit as an expression product of the DNA, and collecting the produced β subunit.

(8) The DNA according to item (3) or (4), further including the nucleotide sequence encoding an α subunit and a γ subunit of glucose dehydrogenase of *Burkholderia cepacia*.

(9) A recombinant vector including the DNA according to item (8).

(10) A transformant transformed with the DNA according to item (8) or the recombinant vector according to item (9).

(11) A method of producing a glucose dehydrogenase complex, including culturing the transformant according to item (10) to produce a glucose dehydrogenase complex as an expression product of the DNA, and collecting the produced complex.

Best Mode for Carrying out the Invention

[0013] Hereinafter, the present invention will be described in detail.

[0014] The inventors of the present invention have searched and isolated a DNA encoding a GDH β subunit of *Burkholderia cepacia* KS1 strain. The aforementioned strain was deposited at International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (Central 6, 1-1-1, Higashi 1-chome, Tsukuba-shi, Ibaraki-ken, Japan, postal code: 305-8566) on September 25, 2000 and received a microorganism accession number of FERM BP-7306. In the present specification, the DNA encoding the GDH β subunit is sometimes referred to as the DNA of the present invention, " β subunit structural gene", or simply " β subunit gene".

[0015] The inventors of the present invention have confirmed that GDH produced by *Burkholderia cepacia* KS1 strain is a polymeric protein containing an α subunit, a β subunit, and a γ subunit. The protein of the present invention corresponds to the β subunit out of these subunits. Spectrophotometric analyses for GDH indicate that the absorption wavelength of an oxidized GDH resembles the absorption wavelengths of alcohol dehydrogenase and aldehyde dehydrogenase of *Gluconobacter* sp. and *Acetabacter* sp. that are composed of dehydrogenase cytochrome complex, and this absorption is lost by heat treatment. This fact and the difference in optimum reaction temperature of the GDH between presence and absence of the β subunit as described below have suggested that the β subunit is composed of cytochrome C.

[0016] Physical and chemical properties of the above GDH are shown below.

(1) Function: the enzyme catalyzes dehydrogenation reaction of glucose.

EP 1 498 484 A1

(2) The enzyme consists of subunits showing a molecular weight of about 60 kDa and a molecular weight of about 43 kDa in SDS-polyacrylamide gel electrophoresis under a reducing condition.

(3) The enzyme shows a molecular weight of about 380 kDa in gel filtration chromatography using TSK Gel G3000SW (Manufactured by Tosoh Corporation).

(4) Optimal reaction temperature: around 45°C (Tris-HCl buffer, pH 8.0).

Physical and chemical properties of an α -subunit alone are shown below.

(1) The protein has a glucose dehydrogenase activity.

(2) The protein shows a molecular weight of about 60 kDa in SDS-polyacrylamide gel electrophoresis under a reducing condition.

(3) Optimal reaction temperature: around 75°C (Tris-HCl buffer, pH 8.0).

[0017] The β subunit can be obtained together with other subunits from a culture of *Burkholderia cepacia* KS1 strain by purifying a GDH complex using GDH activity as an index. The GDH activity can be measured in the same manner as in the known GDH activity measurement. Specifically, the measurement can be performed as follows. A 10 mM potassium phosphate buffer (pH 7.0) containing 594 μ M 1-methoxyphenazine methosulfate (mPMS) and 5.94 μ M 2,6-dichlorophenol indophenol (DCIP) are added to an enzyme sample and glucose as a substrate, and the mixture is incubated at 37°C. A change in the absorbance of DCIP at 600 nm is traced using a spectrophotometer, and the decrease rate in the absorbance is defined as an enzyme reaction rate.

[0018] In addition, since the nucleotide sequence of a gene encoding the β subunit (SEQ ID NO: 15) has been determined by the present invention, the β subunit can also be produced by expressing a DNA having the nucleotide sequence or a DNA encoding the same amino acid sequence as the amino acid sequence encoded by this DNA in a suitable host. The amino acid sequence that can be encoded by the open reading frame (ORF) of SEQ ID NO: 15 is shown in SEQ ID NO: 16. The N-terminal amino acid sequence of the β subunit determined from the protein was identical to the amino acids 23 to 38 of SEQ ID NO: 16. Therefore, the amino acids 1 to 22 are presumed to be a signal peptide. Note that although amino acid residue 1 is described as Val in SEQ ID NOs: 15 and 16, it has a high possibility to be Met and it also has a possibility to be dropped after the translation.

[0019] Results of homology search on the aforementioned amino acid sequence by BLAST indicated overall high homologies: 65% homology with a cytochrome C subunit of oxidoreductase dehydrogenase derived from *Ralstonia solanacearum*, 48% homology with a cytochrome C subunit of sorbitol dehydrogenase derived from *Gluconobacter oxydans*, 44% homology with a cytochrome C subunit of gluconic acid dehydrogenase derived from *Eriwinia cypripedii*, and 55.7% homology on the nucleotide sequence level or a 46.4% homology on the amino acid level with a cytochrome C subunit of 2-keto-gluconic acid dehydrogenase derived from *Pantoea citrea*. Furthermore, the amino acid sequences of these cytochromes C reserved a hem-linking motif (SEQ ID NO: 18). These facts show that the β subunit of the present invention is cytochrome C.

[0020] The β subunit of the present invention may be a protein having the amino acid sequence consisting of the amino acids 23 to 425 of SEQ ID NO: 16 including substitution, deletion, insertion, or addition of 1 to 20, preferably 1 to 10, more preferably 1 to 5 amino acid residues in the amino acid sequence so long as it can function as a GDH β subunit. Note that the term "functions as a GDH β subunit" means "functions as cytochrome C without deteriorating the enzyme activity of the GDH".

[0021] The DNA of the present invention is a DNA that encodes the aforementioned β subunit and can be obtained from, for example, *Burkholderia cepacia* KS1 strain. The DNA of the present invention has been isolated from the chromosomal DNA of *Burkholderia cepacia* KS1 strain in the course of completion of the present invention. The DNA of the present invention can be obtained, for example, by PCR using primers having nucleotide sequences of SEQ ID NOs: 13 and 14 and the chromosomal DNA of *Burkholderia cepacia* KS1 strain as a template. In addition, since the nucleotide sequence of the DNA of the present invention and the amino acid sequence encoded by the nucleotide sequence have been clarified by the present invention, the DNA of the present invention can also be obtained by performing chemical synthesis based on these sequences. Furthermore, the DNA of the present invention can be obtained from the chromosomal DNA of *Burkholderia cepacia* KS1 strain by hybridization using the oligonucleotide prepared based on the aforementioned sequences as probes. Similarly, variants can be obtained from strains other than *Burkholderia cepacia*.

[0022] The DNA of the present invention may be one encoding a protein having the amino acid sequence consisting of the amino acids 23 to 425 of SEQ ID NO: 16 or one having the amino acid sequence including substitution, deletion, insertion, or addition of 1 to 20, preferably 1 to 10, more preferably 1 to 5 amino acid residues and encoding a protein that functions as a GDH β subunit.

[0023] Specifically, the DNA of the present invention includes a DNA having the nucleotide sequence consisting of the nucleotides 187 to 1398 of SEQ ID NO: 15. Further, the DNA of the present invention may be a DNA that hybridizes with SEQ ID NO: 15 or a probe that can be prepared from this sequence under stringent conditions and encodes a

EP 1 498 484 A1

protein that can function as a β subunit. The stringent conditions include those conditions whereby DNAs having a 70% or more, preferably 80% or more, more preferably 90% or more homology to each other hybridize, specifically conditions of 1 x SSC, 0.1% SDS, and 60°C.

[0024] The GDH β subunit can be produced by culturing a transformant that harbors the DNA of the present invention or a recombinant vector containing the DNA of the present invention to produce the GDH β subunit as an expression product of the DNA, and by collecting the GDH β subunit from the microorganism cells or the culture medium. In this case, the DNA encoding the GDH β subunit of the present invention may be expressed together with a DNA encoding an α subunit or further a DNA encoding a γ subunit to produce a GDH complex. A DNA fragment that sequentially encodes the γ subunit and the α subunit can be obtained by PCR using primers having nucleotide sequences of SEQ

ID NOs: 18 and 19.

[0025] Examples of the microorganism that produces the GDH β subunit or the GDH complex include: enterobacteria including *Escherichia coli*; Gram negative bacteria such as *Pseudomonas* and *Gluconobacter*; Gram positive bacteria including bacteria belonging to the genus *Bacillus* such as *Bacillus subtilis*; yeasts such as *Saccharomyces cerevisiae*; and molds such as *Aspergillus niger*. However, the microorganism is not limited to these and any microorganism may be used so long as it is a host microorganism suitable for the production of foreign proteins.

[0026] Vectors that are used for cloning or expressing the DNA of the present invention are suitably those constructed for gene recombination from plasmids or phages that can autonomously replicate in host microorganisms. Examples of vectors for *E. coli* include pBR322, pUC18, pUC118, pUC19, pUC119, pTrc99A, pBluescript, or SuperCos1, which is a cosmid. Transfer of the DNA from the vector that has been used in cloning the DNA of the present invention to other recombinant vectors suitable for expression, etc. can be readily performed by recovering the DNA of the present invention from a recombinant vector containing the DNA of the present invention with a restriction enzyme or by the PCR method and ligating it with a vector fragment. Furthermore, transformation of microorganisms with these vectors can be performed by the competent cell method by treatment with calcium for bacteria belonging to the genus *Escherichia*, the protoplast method for bacteria belonging to the genus *Bacillus*, the KU method or the KUR method for yeasts, and the micromanipulation method for molds and so forth. In addition, the electroporation method can also be used widely.

[0027] Selection of host microorganisms based on presence or absence of introduction of the target recombinant vector therein may be performed by using a chemical resistance marker of the vector containing the target DNA and the like. For example, a microorganism that can grow in a selective medium based on a chemical resistance marker and produces GDH may be selected.

[0028] As for the culture method of the transformant, culture conditions may be selected by considering nutritional and physiological properties of the host. In many cases, liquid culture is performed. It is industrially advantageous to perform aeration culture with shaking.

[0029] As nutrients of the medium, those usually used for culture of microorganisms may be widely used. As carbon sources, any assimilable carbon compounds may be used, and examples of the compounds to be used include glucose, sucrose, lactose, maltose, lactose, molasses, pyruvic acid and so forth. Furthermore, as nitrogen sources, any utilizable nitrogen compounds may be used, and examples of the compounds to be used include peptone, meat extracts, yeast extracts, casein hydrolysates, soybean cake alkaline extracts and so forth. In addition, phosphate, carbonate, sulfate, salts of magnesium, calcium, potassium, iron, manganese, zinc and so forth, particular amino acids, particular vitamins and so forth are used as required.

[0030] Although the culture temperature can be appropriately changed in a range in which a bacteria grows and produces the protein of the present invention, it is preferably about 20°C to 42°C. The culture time somewhat varies depending on the conditions. However, the culture may be completed at an appropriate time estimated to give the maximum GDH yield, and the culture time is usually about 12 to 72 hours. Although pH of the medium may be appropriately changed in a range in which a bacteria grows and produces the protein of the present invention, it is preferably in the range of about pH 6.0 to 9.0.

[0031] The culture medium containing cells of the microorganism producing the protein of the present invention in the culture can be collected and utilized as they are. However, when the protein of the present invention exists in the culture medium, the culture medium is usually separated into the solution containing the protein of the present invention and microorganism cells by filtration, centrifugation or the like in a conventional manner and then used. When the protein of the present invention exists in the cells, the cells are collected from the obtained culture by means of filtration, centrifugation or the like, and then disrupted by a mechanical method or an enzymatic method such as use of lysozyme. Further, a chelating agent such as EDTA and a surfactant are added to the cell to solubilize the protein of the present invention, as required, followed by isolation and collection as an aqueous solution.

[0032] Protein may be precipitated from the thus-obtained protein-containing solution by, for example, vacuum concentration, membrane concentration, salting out with ammonium sulfate, sodium sulfate or the like, or a fractional precipitation with a hydrophilic organic solvent such as methanol, ethanol, and acetone. Further, heat treatment and isoelectric point treatment are also effective purification means. Then, purification can be performed by a suitable

EP 1 498 484 A1

combination of gel filtration using an adsorbent, gel filtration agent, etc., absorption chromatography, ion exchange chromatography and affinity chromatography to obtain a purified protein of the present invention.

[0033] A purified enzyme preparation can be obtained by isolation and purification based on column chromatography. Although the purified enzyme preparation is preferably purified to such an extent that a single band is obtained in electrophoresis (SDS-PAGE), it may contain an α -subunit or a γ -subunit.

[0034] The thus-obtained purified enzyme can be powdered by, for example, lyophilization, vacuum drying, spray drying or the like and distributed.

[0035] The GDH complex consisting of a β -subunit, an α -subunit, or whenever required, a γ -subunit of the present invention, or transformants containing those, may be used for an enzyme electrode of a glucose sensor. As the electrode, a carbon electrode, gold electrode, platinum electrode or the like may be used, and the GDH of the present invention is immobilized on the electrode. Examples of the method for immobilization include a method of using a crosslinking reagent, a method of entrapping the enzyme in a polymer matrix, a method of covering the enzyme with a dialysis membrane, methods of using a photocrosslinking polymer, conductive polymer, oxidation-reduction polymer or the like. Alternatively, the enzyme may be immobilized in a polymer or immobilized on an electrode by adsorption together with an electronic mediator of which typical examples are ferrocene and derivatives thereof, or these methods may be used in combination. Typically, the glucose dehydrogenase of the present invention is immobilized on a carbon electrode by using glutaraldehyde, and then glutaraldehyde is blocked by a treatment with a reagent having an amine group.

[0036] The glucose concentration can be measured as follows. A buffer is placed in a thermostatic cell, and a mediator is added thereto. Then, a constant temperature is maintained. As the mediator, potassium ferricyanide, phenazine methosulfate and so forth may be used. An electrode on which the enzyme of the present invention is immobilized is used as a working electrode, and a counter electrode (e.g., platinum electrode) and a reference electrode (e.g., Ag/AgCl electrode) are used. After a constant voltage is applied to the carbon electrode to obtain a steady-state current, a sample containing glucose is added thereto and the increase of the current is measured. The glucose concentration in the sample can be calculated according to a calibration curve produced by using glucose solutions having standard concentrations.

[0037] The GDH complex containing the β subunit of the present invention may be used as a component in an assay kit for saccharides such as glucose. Typically, the kit includes in addition to the GDH complex, a buffer necessary for the assay, a mediator, a standard solution of, for example, glucose for preparing a calibration curve, and a manual on the use of the kit. The enzyme according to the present invention may be provided in various forms, for example, as a freeze-dried reagent or as a solution in a suitable stock solution.

Example

[0038] Hereinafter, the present invention will be described specifically with reference to examples.

Reference Example 1: Isolation of a gene encoding GDH α subunit of *Burkholderia cepacia* KS1 strain

<1> Preparation of chromosomal DNA from *Burkholderia cepacia* KS1 strain.

[0039] A chromosomal gene was prepared from the *Burkholderia cepacia* KS1 strain in a conventional manner. That is, the bacterial strain was shaken overnight at 34°C by using a TL liquid medium (10 g of polypeptone, 1 g of yeast extract, 5 g of NaCl, 2 g of KH_2PO_4 , 5 g of glucose in 1 L, pH 7.2). The grown cells were collected by centrifugation. The cells were suspended in a solution containing 10 mM NaCl, 20 mM Tris-HCl (pH 8.0), 1 mM EDTA, 0.5% SDS, and 100 $\mu\text{g}/\text{ml}$ proteinase K and treated at 54°C for 6 hours. This mixture was added to an equivalent volume of phenol-chloroform and stirred at room temperature for 10 minutes, and then the supernatant was collected by centrifugation. The supernatant was added to sodium acetate to give a final concentration of 0.3 M and overlaid with two-fold volume of ethanol to precipitate chromosomal DNA in the intermediate layer. The DNA was taken up with a glass rod, washed with 70% ethanol, and dissolved in an appropriate amount of TE buffer to obtain a chromosomal DNA solution.

<2> Determination of N-terminus amino acid sequence of GDH α -subunit

[0040] GDH purified in the same manner as in Example 2 was concentrated by lyophilization and separated by SDS-electrophoresis using 12.5% polyacrylamide to isolate the α -subunit. The thus obtained α -subunit was transferred onto a polyvinylidene fluoride membrane, and then the N-terminus amino acid sequence was determined by using an amino acid sequencer (PPSQ-10, manufactured by Shimadzu Corporation). As a result, it was found that the enzyme contained a peptide sequence consisting of 11 residues of the amino acid numbers 2 to 12 in the amino acid sequence of SEQ ID NO: 3.

EP 1 498 484 A1

<3> Cloning of a gene encoding α -subunit

[0041] The DNA prepared in <1> (1 μ g) was subjected to partial digestion with a restriction enzyme *Sau3AI*, followed by treatment with calf intestinal alkaline phosphatase (CIAP). Separately, SuperCosI (obtained from Stratagene), which is a cosmid, was treated with *Bam*HI, and the DNA fragment obtained by the partial digestion of the chromosomal DNA fragment derived from the α -15 strain with *Sau3AI* was introduced into SuperCosI by using T4 DNA ligase. *E. coli* XL-1 Blue MR (obtained from Stratagene) was transformed with the obtained recombinant DNA. A transformant was selected on an LB agar medium containing 10 μ g/ml neomycin and 25 μ g/ml ampicillin based on neomycin resistance and ampicillin resistance, which are antibiotic resistances on SuperCosI. The obtained transformant was cultured in the LB liquid medium. These transformant cells were collected and suspended in a reagent for measuring the GDH activity, and a clone was selected by using dehydrogenase activity for glucose as an index. As a result, one clone showing the glucose dehydrogenase activity was obtained.

<4> Subcloning

[0042] DNA fragments containing the target gene were prepared from the cosmid, SuperCosI, containing the gene encoding the α -subunit obtained in <3>. The inserted gene fragments were cleaved from the cosmid by using a restriction enzyme *Nof*I. These DNA fragments were treated with a restriction enzyme *Xba*I and introduced into a plasmid pUC18 digested with *Xba*I. The *E. coli* DH5 α -MCR strain was transformed with the plasmid pUC18 containing each inserted fragment, and colonies appeared on an LB agar medium containing 50 μ g/ml of ampicillin were collected. The obtained transformants were cultured in the liquid LB medium, followed by examination for the GDH activity in the cells in the same manner as in <3>. As a result, a transformant showing the GDH activity was obtained. The plasmid was extracted from this transformant, and the inserted DNA fragment was analyzed. As a result, an insert of about 8.8 kbp was confirmed. This plasmid was designated as pKS1.

<5> Determination of the nucleotide sequence

[0043] Restriction enzyme analysis of the inserted DNA fragment in pKS1 was performed and the nucleotide sequence of the fragment was determined according to the conventional method. As a result, the sequence of the DNA encoding the N-terminus amino acid sequence of the α -subunit found in <2> was confirmed in this inserted DNA fragment, and an open reading frame containing this sequence was found. The determined nucleotide sequence and the amino acid sequence that can be encoded by this nucleotide sequence are as shown in SEQ ID NOs: 1 and 3. In the nucleotide sequence of SEQ NO: 1, nucleotide sequence downstream from nucleotide number 2,386 encode amino acid sequence of SEQ NO: 4, and encode β -subunit.

Reference Example 2: Production of a GDH- α -subunit by recombinant *E. coli*.

[0044] Since the nucleotide sequence of the α -subunit was determined, a vector was prepared using the aforementioned structural gene of the α -subunit, and a transformant was further produced using this vector.

[0045] First, a gene to be inserted into the vector was prepared as follows.

[0046] Amplification was performed by PCR using a genome fragment derived from the KS1 strain as a template so that a desired restriction enzyme site is included. The following pair of oligonucleotide primers were used in PCR.

(Forward)

5' - CCCAAGCTTGGGCCGATACCGATACGCA - 3' (SEQ ID NO: 5)

(Reverse)

5' - GAGAAGCTTTCGCGACGGTCAGACTTCC - 3' (SEQ ID NO: 6)

[0047] The gene amplified by PCR was digested with a restriction enzyme *Hind*III and inserted into an expression vector pFLAG-CTS (SIGMA) at its cloning site, *Hind*III site. The obtained plasmid was designated as pFLAG-CTS/ α .

[0048] The *E. coli* DH5 α MCR strain was transformed with the aforementioned plasmid pFLAG-CTS/ α , and colonies appeared on the LB agar medium containing 50 μ g/ml of ampicillin were collected.

[0049] Further, when the open reading frame of the pKS1 inserted fragment was searched in the upstream of the α -subunit, a structural gene consisting of 507 nucleotides encoding a polypeptide including 168 amino acid residues

EP 1 498 484 A1

shown in SEQ ID NO: 2 (nucleotide numbers 258 to 761 in SEQ ID NO: 1) was newly found. This structural gene was considered to encode the γ -subunit.

[0050] Since it was found that the region encoding the γ -subunit existed upstream of the coding region of the α -subunit, a recombinant vector containing a gene having a polycistronic structure continuously including the γ -subunit and the α -subunit was produced, and a transformant introduced with this vector was constructed.

[0051] First, a gene to be inserted into the vector was prepared as follows.

[0052] Amplification was performed by PCR using a genome fragment derived from the KS1 strain continuously including the structural gene of the γ -subunit and the structural gene of the α -subunit as a template so that a desired restriction enzyme site is included. The following pair of oligonucleotide primers were used for PCR.

(Forward)

5'-CATGCCATGGCACACAACGACAACACT-3' (SEQ ID NO: 7)

(Reverse)

5'-CCCAAGCTTGGGTCAGACTTCCTTCTCAGC-3' (SEQ ID NO: 8)

[0053] The 5'-terminus and the 3'-terminus of the gene amplified by PCR were digested with *Nco*I and *Hind*III, respectively, and the gene was inserted into the vector pTrc99A (Pharmacia) at its cloning site, *Nco*I/*Hind*III site. The obtained plasmid was named pTrc99A/ γ - α .

[0054] The *E. coli* DH5 α MCR strain was transformed with the aforementioned plasmid pTrc99A/ γ - α , and colonies appeared on the LB agar medium containing 50 μ g/ml of ampicillin were collected.

[0055] The α -subunit was produced using the *E. coli* DH5 α MCR strain transformed with each of the aforementioned plasmids pKS1, pFLAG-CTS/ α and pTrc99A/ γ - α . Each transformant was inoculated into 3 ml of the LB medium containing 50 μ g/ml of ampicillin and cultured at 37°C for 12 hours, and the cells were collected by centrifugation. The cells were disrupted by using a French press (1,500 kgf), and a membrane fraction (10 mM potassium phosphate buffer, pH 6.0) was isolated by ultracentrifugation (4°C, 160,400 x g, 90 minutes).

Reference Example 3: Confirmation of GDH activity

[0056] First, the GDH activity in each of the aforementioned membrane fractions was confirmed. Specifically, visual determination was performed by using 10 mM potassium phosphate buffer (pH 7.0) containing 594 μ M of methylphenazine methosulfate (mPMS) and 5.94 μ M of 2,6-dichlorophenolindophenol (DCIP). The results are shown below. The number of + represents the degree of color change from blue to colorless.

Membrane fraction of cultured transformant transformed with pFLAG-CTS/ α : +

Membrane fraction of cultured transformant transformed with pKS1 : ++

Membrane fraction of cultured transformant transformed with pTrc99A/ γ - α : +++

[0057] The GDH activity of the membrane fraction of the cultured transformant transformed with pFLAG-TS/ α containing only with the α -subunit was the lowest, and the GDH activity of the membrane fraction of the cultured transformant transformed with pTrc99A/ γ - α , with which a vector was efficiently constructed, was the highest.

[0058] Although the α -subunit was expressed even in the transformant transformed with a vector using only the structural gene of the α -subunit, the α -subunit could be efficiently obtained by using a vector containing the structural gene of the γ -subunit and the structural gene of the α -subunit in combination.

[0059] Glucose was assayed using the glucose dehydrogenase of the present invention. The enzymatic activity of the glucose dehydrogenase (α -subunit) of the present invention was measured by using glucose having various concentrations. The GDH activity was measured in 10 mM potassium phosphate buffer (pH 7.0) containing 594 μ M of methylphenazine methosulfate (mPMS) and 5.94 μ M of 2,6-dichlorophenolindophenol (DCIP). An enzyme sample and glucose serving as a substrate were added thereto, followed by incubation at 37°C, and change in the absorbance of DCIP at 600nm was monitored by using a spectrophotometer. The decreasing rate of the absorbance was measured as an enzymatic reaction rate. Glucose could be quantified in the range of 0.01 to 1.0 mM using the GDH of the present invention.

EP 1 498 484 A1

Example 1: Isolation of gene encoding a GDH β -subunit of *Burkholderia cepacia* KS1 strain

<1> Searching for *Burkholderia cepacia* KS1 strain GDH β subunit

- 5 [0060] GDH β subunit gene derived from KS1 strain was searched using *Burkholderia cepacia* J2315 strain genome database of Sanger Centre (<http://www.sanger.ac.uk/>). Referring to a known N-terminal sequence of KS1 strain GDH β subunit (SEQ ID NO: 9), there was designed an amino acid sequence (SEQ ID NO: 10) that has high homology to each cytochrome c subunit of an alcohol dehydrogenase derived from *Acetobacter* Sp. or *Gluconobacter* Sp. (Tamaki T. et al., *Biochim Biophys Acta* 1088 (2): 292-300 (1991), Matsushita K., et al., *Biosci. Biotech. Biochem.*, 56, 304-310 (1992), Takemura H., et al., *J Bacteriol.* 175, 6857-66 (1993), Kondo K. et al., *Appl Environ Microbiol.* 63, 1131-8 (1997)); a gluconate dehydrogenase derived from *Erwinia* sp. or *Pseudomonas* sp. (Yum DY, et al., *J Bacteriol.* 179, 6566-72 (1997), Matsushita K. et al., *J Biochem.* 85, 1173-81 (1979)); a sorbitol dehydrogenase derived from *Gluconobacter* sp. (Choi, E.S., et al., *FEMS Microbiol. Lett.*, 125, 45-50 (1995)); and a 2-ketogluconate dehydrogenase derived from *Erwinia* sp. or *Pantoea* sp. (Pujol CJ et al., *J Bacteriol.* 182, 2230-7, (2000)).
- 10 [0061] Based on the aforementioned amino acid sequence, gene sequences that encode amino acid sequences having high homologies have been searched from the aforementioned database of *Burkholderia cepacia* J2315 strain by BLAST. Then, the obtained five sequences were searched for homology with the C-terminal sequence of the GDH α subunit of KS1 strain. As a result, amino acid sequences translated from two gene fragments showed high homologies (>90%). Each gene fragment was as short as 200 to 500 bp, so that sequences having high homologies with these sequences were searched from the genome database of *Burkholderia cepacia* J2315 strain by BLAST and the fragments were joined each other. As a result, a fragment of 3,110 bp was obtained. In the obtained nucleotide sequence, there existed an ORF that is presumed to be the C-terminus of the GDH and an ORF that is presumed to be cytochrome C structural gene of 1,275 bp (SEQ ID NO: 11). The amino acid sequence encoded by the ORF is shown in SEQ ID NO: 12. Results of comparison between the obtained nucleotide sequence of the J2315 strain and the nucleotide sequence of the α subunit of KS1 strain that has already been cloned indicate that in the downside of the α subunit, the nucleotide sequence having a high homology with the nucleotide sequence encoding the signal peptide of cytochrome C of J2315 strain is contained.
- 15 [0062] From the above, the third ORF in the cloned fragment of *Burkholderia cepacia* KS1 strain obtained in Reference Example 1 (nucleotides 2386 et seq. of SEQ ID NO: 1) is presumed to encode the β subunit. The amino acid sequence at the N-terminus of the purified β subunit corresponds to the 5 amino acid residues translated by the nucleotide sequence consisting of nucleotides 2452 to 2466 in SEQ ID NO: 1, which also suggests that the aforementioned ORF encodes the β subunit.
- 20
- 25

<2> Amplification of β subunit structural gene using inverse PCR method

(1) Culture of microorganism cell and extraction of genome

35 [0063] Using 5 ml of complete medium (0.5% polypepton, 0.3% yeast extract, 0.5% NaCl), KS1 strain was cultured with shaking at 37°C overnight. Genome was extracted from the obtained microorganism cells using GennomicPrep™ Cells and Tissue DNA Isolation Kit (Amersham Pharmacia Biotech). The method was performed in accordance with the attached manual. The obtained genome was subjected to phenol/chloroform treatment and precipitated with ethanol, and then dissolved in purified water.

40

(2) Cyclization of genome fragment

45 [0054] The genome extracted from the KS1 strain was digested with *Bam*HI, *Eco*RI, *Hind*III, *Sma*I, *Sac*I, and *Xho*I and the genome fragments were recovered by precipitation with ethanol. Then, 1 μ g of genome digested with the restriction enzymes was subjected to ligation reaction using a DNA ligation kit (Takara Shuzo Co., Ltd.) at 16°C overnight.

50

(3) PCR

55 [0065] 50 pmol of forward primer (EF1 SEQ ID NO: 13) designed based on the nucleotide sequence in the N-terminal signal sequence region of the GDH β subunit of the KS1 strain, 50 pmol of reverse primer (ER1 SEQ ID NO: 14) (all the primers were synthesized by Invitrogen on consignment), 0.5 ml of LATAq (Takara Bio Co., Ltd.), 8 μ l of dNTP solution, and 5 μ l of 10 \times PCR buffer was added to purified water so as to have a total volume of 50 μ l, and PCR was performed using a program temp control system PC-801 (ASTECH). The PCR reaction was performed under the following conditions: after 30 cycles of 94°C for 5 minutes, 98°C for 20 seconds, 62°C for 30 seconds, 72°C for 6 minutes,

EP 1 498 484 A1

and 72°C for 10 minutes.

[0066] When the genome digested with a restriction enzyme (*Sma*) is used as a template, a fragment having a size of about 2.1 kbp was confirmed on Agarose electrophoresis.

5. <3> Sequencing of PCR-amplified fragment

(1) TA cloning

[0067] After the aforementioned inverse PCR product was electrophoresed on Agarose gel, the band was cut out and purified using Gene clean II KIT (Bio101 inc.). The fragment was ligated to pGEM-T Vector using pGEM-T and pGEM-T EASY Vector Systems (Promega). *E. coli* DH5 α was transformed with the ligated vector, and the transformant was cultured on an L-agar medium containing 50 μ g/ml ampicillin, 40 μ g/ml X-Gal, and 0.1 μ M IPTG overnight. From the appeared colonies, white colonies were selected and cultured in an L medium containing 50 μ g/ml ampicillin overnight, followed by extraction of plasmids from the cells by the alkali method.

(2) Preparation of sequence sample

[0068] The obtained plasmid was treated with RNase and 0.6 volume of 20% PEG6000/2.5 M NaCl was added thereto. The mixture was left to stand on ice for 1 hour. Thereafter, the mixture was centrifuged at 15,000 rpm and 4°C for 15 minutes to obtain a pellet. The pellet was washed with 70% ethanol and dried in vacuum. The dried product was dissolved in purified water.

(3) Analysis of nucleotide sequence of DNA

[0069] The nucleotide sequence of the inserted fragment of the plasmid obtained in (2) was analyzed using ABI PRISM™ 310 Genetic Analyzer (PERKIN-ELMER Applied Biosystems). A portion of the sequence of the inserted fragment was determined from the multicloning site of the vector using M13 primer. As a result, the nucleotide sequence containing the N-terminus of the β subunit that had been already analyzed was confirmed. Based on this sequence, primers were sequentially prepared and used to determine the nucleotide sequence of the inserted fragment. The result is shown in SEQ ID NO: 15. Further, the amino acid sequence encoded by the ORF in the nucleotide sequence is shown in SEQ ID NO: 16.

[0070] The β subunit has 425 amino acid residues in total, and from the comparison with the N-terminal amino acid sequence already obtained, 22 residues among them are considered to be a signal peptide. The molecular weight of the β subunit calculated based on the amino acid sequence is 45,276 Da and the molecular weight 42,731 Da of the portion excluding the signal peptide is substantially identical to the molecular weight 43 kDa of the GDH β subunit of the KS1 strain. In the amino acid sequence of the β subunit, linking motifs (SEQ ID NO: 18) that links with hem in cytochrome C were confirmed at 3 positions. The ORFs were located immediately downstream of the ORF of the structural gene of the α subunit, and a sequence that is presumed to be an SD sequence existed upstream of the initiation codon.

[0071] Homology search for the obtained amino acid sequence by BLAST showed overall high homologies; a 65% homology with cytochrome C subunit of oxidoreductase dehydrogenase derived from *Ralstonia solanacearum*, a 48% homology with a cytochrome C subunit of sorbitol dehydrogenase derived from *Gluconobacter oxydans*, a 44% homology with a cytochrome C subunit of gluconic acid dehydrogenase derived from *Eriwinia cypripedii*, and a 46.4% homology on an amino acid level with a cytochrome C subunit of 2-keto-gluconic acid dehydrogenase derived from *Pantoea citrea*. Furthermore, the amino acid sequences of cytochromes C reserved a hem-linking motif (SEQ ID NO: 18).

[0072] The structural gene of the GDH β subunit of the KS1 strain has a homology of 92.0% on the nucleotide sequence level and of 92.2% on an amino acid level with the structural gene of the GDH β subunit of the J2315 strain.

50 Industrial Applicability

[0073] The present invention provides the GDH β subunit of a microorganism belonging to the genus *Burkholderia* and the DNA encoding it.

55

EP 1 498 484 A1

SEQUENCE LISTING

5 <110> SODE, Koji

<120> Glucose dehydrogenase beta-subunit and DNA encoding the same

10 <130> G780-OP1551

<141> 2003-04-25

15 <150> JP 2002-125353

<151> 2002-04-26

<160> 19

20 <170> PatentIn Ver. 2.0

<210> 1

25 <211> 2467

<212> DNA

<213> Burkholderia cepacia

30 <220>

<221> CDS

<222> (258)..(761)

35 <220>

<221> CDS

<222> (764)..(2380)

40 <220>

<221> CDS

<222> (2386)..(2466)

45 <400> 1

aagcttctcgt ttgatctga cgcgattcta accgagcgtc tglgaggcgg aacgcgacat 60

gcttcgtgtc gcacacggtt cgcgccgacg acacaaaaat gcagcgaat ggcgtgacgt 120

tacgaalggt tgacacatig aalggaciat aaaaccatig tccgttccgg aatgtgcggt 180

50 tacaattcag gtccgcgcgg atttttgaga aataicaagg ggggttttcc cgaatccgg! 240

gttcgagaga aggaaac atg cac aac gac aac act ccc cac tgg cgt cgc 290

Met His Asn Asp Asn Thr Pro His Ser Arg Arg

1 5 10

55

EP 1 498 484 A1

Ser Ser Pro Trp Ala Pro His Pro Glu Tyr Gly Pro Pro Asn Asp Tyr
 235 240 245 250
 5 ctc atc ctg aag ggc gag cac aag ttc aac tgc cag tac atc cgc gcg 1057
 Leu Ile Leu Lys Gly Glu His Lys Phe Asn Ser Gln Tyr Ile Arg Ala
 255 260 265
 10 gtg ggc ggc acg acg tgg cac tgg gcc gcg tgc gcg tgg cgc ttc att 1105
 Val Gly Gly Thr Thr Trp His Trp Ala Ala Ser Ala Trp Arg Phe Ile
 270 275 280
 ccg aac gac ttc aag atg aag agc gtg tac ggc gtc ggc cgc gac tgg 1153
 Pro Asn Asp Phe Lys Met Lys Ser Val Tyr Gly Val Gly Arg Asp Trp
 285 290 295
 15 ccg atc cag tac gac gat ctc gag ccg tac tat cag cgc gcg gag gaa 1201
 Pro Ile Gln Tyr Asp Asp Leu Glu Pro Tyr Tyr Gln Arg Ala Glu Glu
 300 305 310
 20 gag ctc ggc gtg tgg ggc ccg ggc ccc gag gaa gat ctg tac tgc ccg 1249
 Glu Leu Gly Val Trp Gly Pro Gly Pro Glu Glu Asp Leu Tyr Ser Pro
 315 320 325 330
 25 cgc aag cag ccg tat ccg atg ccg ccg ctg ccg ttg tgc ttc aac gag 1297
 Arg Lys Gln Pro Tyr Pro Met Pro Pro Leu Pro Leu Ser Phe Asn Glu
 335 340 345
 cag acc atc aag acg gcg ctg aac aac tac gat ccg aag ttc cat gtc 1345
 Gln Thr Ile Lys Thr Ala Leu Asn Asn Tyr Asp Pro Lys Phe His Val
 350 355 360
 30 gtg acc gag ccg gtc gcg cgc aac agc cgc ccg tac gac ggc cgc ccg 1393
 Val Thr Glu Pro Val Ala Arg Asn Ser Arg Pro Tyr Asp Gly Arg Pro
 365 370 375
 35 act tgt tgc ggc aac aac aac tgc atg ccg atc tgc ccg atc ggc gcg 1441
 Thr Cys Cys Gly Asn Asn Asn Cys Met Pro Ile Cys Pro Ile Gly Ala
 380 385 390
 40 atg tac aac ggc atc glg cac gtc gag aag gcc gaa cgc gcc ggc gcg 1489
 Met Tyr Asn Gly Ile Val His Val Glu Lys Ala Glu Arg Ala Gly Ala
 395 400 405 410
 aag ctg atc gag aac gcg gtc gtc tac aag ctg gag acg ggc ccg gac 1537
 Lys Leu Ile Glu Asn Ala Val Val Tyr Lys Leu Glu Thr Gly Pro Asp
 415 420 425
 45 aag cgc atc gtc gcg gcg ctc tac aag gac aag acg ggc gcc gag cat 1585
 Lys Arg Ile Val Ala Ala Leu Tyr Lys Asp Lys Thr Gly Ala Glu His
 430 435 440
 50 cgc gtc gaa ggc aag tat ttc gtg ctc gcc gcg aac ggc atc gag acg 1633
 Arg Val Glu Gly Lys Tyr Phe Val Leu Ala Ala Asn Gly Ile Glu Thr
 445 450 455
 55 ccg aag atc ctg ctg atg tcc gcg aac cgc gat ttc ccg aac ggt gtc 1681
 Pro Lys Ile Leu Leu Met Ser Ala Asn Arg Asp Phe Pro Asn Gly Val

EP 1 498 484 A1

460 465 470
 gcg aac agc tgc gac atg gtc ggc cgc aac ctg atg gac cat ccg ggc 1729
 Ala Asn Ser Ser Asp Met Val Gly Arg Asn Leu Met Asp His Pro Gly
 475 480 485 490
 acc ggc gtg tgc ttc tat ggc agc gag aag ctg tgg ccg ggc cgc ggc 1777
 Thr Gly Val Ser Phe Tyr Ala Ser Glu Lys Leu Trp Pro Gly Arg Gly
 495 500 505
 ccg cag gag atg acg tgc ctg atc ggt ttc cgc gac ggt ccg ttc cgc 1825
 Pro Gln Glu Thr Ser Leu Ile Gly Phe Arg Asp Gly Pro Phe Arg
 510 515 520
 gcg acc gaa ggc ggc aag aag atc cac ctg tgc aac ctg tgc cgc atc 1873
 Ala Thr Glu Ala Ala Lys Lys Ile His Leu Ser Asn Leu Ser Arg Ile
 525 530 535
 gac cag gag acg cag aag atc ttc aag gcc ggc aag ctg atg aag ccc 1921
 Asp Gln Glu Thr Gln Lys Ile Phe Lys Ala Gly Lys Leu Met Lys Pro
 540 545 550
 gac gag ctg gac ggc cag atc cgc gac cgt tcc gca cgc tac gtg cag 1969
 Asp Glu Leu Asp Ala Gln Ile Arg Asp Arg Ser Ala Arg Tyr Val Gln
 555 560 565 570
 ttc gac tgc ttc cac gaa atc ctg ccg caa ccc gag aac cgc atc gtg 2017
 Phe Asp Cys Phe His Glu Ile Leu Pro Gln Pro Glu Asn Arg Ile Val
 575 580 585
 ccg agc aag acg ggc acc gat ggc atc ggc att ccg cgc ccc gag atc 2065
 Pro Ser Lys Thr Ala Thr Asp Ala Ile Gly Ile Pro Arg Pro Glu Ile
 590 595 600
 acg tat ggc atc gac gac tac gtc aag cgc ggc gcc ggc cat acg cgc 2113
 Thr Tyr Ala Ile Asp Asp Tyr Val Lys Arg Gly Ala Ala His Thr Arg
 605 610 615
 gag gtc tac ggc acc gcc ggc aag gtc ctg ggc ggc acc gac gtc gtg 2161
 Glu Val Tyr Ala Thr Ala Ala Lys Val Leu Gly Gly Thr Asp Val Val
 620 625 630
 ttc aac gac gaa ttc ggc ccg aac aat cac atc acg ggc tgc acg atc 2209
 Phe Asn Asp Glu Phe Ala Pro Asn Asn His Ile Thr Gly Ser Thr Ile
 635 640 645 650
 atg ggc gcc gat ggc cgc gac tcc gtc gtc gac aag gac tgc cgc acc 2257
 Met Gly Ala Asp Ala Arg Asp Ser Val Val Asp Lys Asp Cys Arg Thr
 655 660 665
 ttc gac cat ccg aac ctg ttc att tgc agc agc ggc acc atg ccg acc 2305
 Phe Asp His Pro Asn Leu Phe Ile Ser Ser Ser Ala Thr Met Pro Thr
 670 675 680
 gtc ggt acc gta aac gtc acg ctg acg atc gcc ggc ctg ggc ctg cgc 2353
 Val Gly Thr Val Asn Val Thr Leu Thr Ile Ala Ala Leu Ala Leu Arg
 685 690 695

EP 1 498 484 A1

atg tgg gac acg clg aag aag gaa gtc tgacc gtc cgg aaa tct act ctc 2403
 Met Ser Asp Thr Leu Lys Lys Glu Val Val Arg Lys Ser Thr Leu
 5 700 705 710
 act ttc ctc atc gcc ggc tgc ctc ggc ttg ccg ggc ttc ggc cgc ggc 2451
 Thr Phe Leu Ile Ala Gly Cys Leu Ala Leu Pro Gly Phe Ala Arg Ala
 715 720 725
 10 gcc gat ggc gcc gat c 2467
 Ala Asp Ala Ala Asp
 730
 15 <210> 2
 <211> 168
 <212> PRT
 <213> Burkholderia cepacia
 20
 <400> 2
 Met His Asn Asp Asn Thr Pro His Ser Arg Arg His Gly Asp Ala Ala
 5 10 15
 25 Ala Ser Gly Ile Thr Arg Arg Gln Trp Leu Gln Gly Ala Leu Ala Leu
 20 25 30
 Thr Ala Ala Gly Leu Thr Gly Ser Leu Thr Leu Arg Ala Leu Ala Asp
 35 40 45
 30 Asn Pro Gly Thr Ala Pro Leu Asp Thr Phe Met Thr Leu Ser Glu Ser
 50 55 60
 Leu Thr Gly Lys Lys Gly Leu Ser Arg Val Ile Gly Glu Arg Leu Leu
 65 70 75 80
 35 Gln Ala Leu Gln Lys Gly Ser Phe Lys Thr Ala Asp Ser Leu Pro Gln
 85 90 95
 Leu Ala Gly Ala Leu Ala Ser Gly Ser Leu Thr Pro Glu Gln Glu Ser
 100 105 110
 40 Leu Ala Leu Thr Ile Leu Glu Ala Trp Tyr Leu Gly Ile Val Asp Asn
 115 120 125
 Val Val Ile Thr Tyr Glu Glu Ala Leu Met Phe Gly Val Val Ser Asp
 130 135 140
 45 Thr Leu Val Ile Arg Ser Tyr Cys Pro Asn Lys Pro Gly Phe Trp Ala
 145 150 155 160
 Asp Lys Pro Ile Glu Arg Gln Ala
 165
 50
 <210> 3
 <211> 539
 <212> PRT
 55 <213> Burkholderia cepacia

EP 1 498 484 A1

<400> 3

5 Met Ala Asp Thr Asp Thr Gln Lys Ala Asp Val Val Val Val Gly Ser
 1 5 10 15
 Gly Val Ala Gly Ala Ile Val Ala His Gln Leu Ala Met Ala Gly Lys
 20 25 30

10 Ala Val Ile Leu Leu Glu Ala Gly Pro Arg Met Pro Arg Trp Glu Ile
 35 40 45
 Val Glu Arg Phe Arg Asn Gln Pro Asp Lys Met Asp Phe Met Ala Pro
 50 55 60

15 Tyr Pro Ser Ser Pro Trp Ala Pro His Pro Glu Tyr Gly Pro Pro Asn
 65 70 75 80
 Asp Tyr Leu Ile Leu Lys Gly Glu His Lys Phe Asn Ser Gln Tyr Ile
 85 90 95

20 Arg Ala Val Gly Gly Thr Thr Trp His Trp Ala Ala Ser Ala Trp Arg
 100 105 110
 Phe Ile Pro Asn Asp Phe Lys Met Lys Ser Val Tyr Gly Val Gly Arg
 115 120 125

25 Asp Trp Pro Ile Gln Tyr Asp Asp Leu Glu Pro Tyr Tyr Gln Arg Ala
 130 135 140
 Glu Glu Glu Leu Gly Val Trp Gly Pro Gly Pro Glu Glu Asp Leu Tyr
 145 150 155 160

30 Ser Pro Arg Lys Gln Pro Tyr Pro Met Pro Pro Leu Pro Leu Ser Phe
 165 170 175
 Asn Glu Gln Thr Ile Lys Thr Ala Leu Asn Asn Tyr Asp Pro Lys Phe
 180 185 190

35 His Val Val Thr Glu Pro Val Ala Arg Asn Ser Arg Pro Tyr Asp Gly
 195 200 205
 Arg Pro Thr Cys Cys Gly Asn Asn Asn Cys Met Pro Ile Cys Pro Ile
 210 215 220

40 Gly Ala Met Tyr Asn Gly Ile Val His Val Glu Lys Ala Glu Arg Ala
 225 230 235 240
 Gly Ala Lys Leu Ile Glu Asn Ala Val Val Tyr Lys Leu Glu Thr Gly
 245 250 255

45 Pro Asp Lys Arg Ile Val Ala Ala Leu Tyr Lys Asp Lys Thr Gly Ala
 260 265 270
 Glu His Arg Val Glu Gly Lys Tyr Phe Val Leu Ala Ala Asn Gly Ile
 275 280 285

50 Glu Thr Pro Lys Ile Leu Leu Met Ser Ala Asn Arg Asp Phe Pro Asn
 290 295 300
 Gly Val Ala Asn Ser Ser Asp Met Val Gly Arg Asn Leu Met Asp His
 305 310 315 320

55 Pro Gly Thr Gly Val Ser Phe Tyr Ala Ser Glu Lys Leu Trp Pro Gly

EP 1 498 484 A1

325 330 335
 Arg Gly Pro Gln Glu Met Thr Ser Leu Ile Gly Phe Arg Asp Gly Pro
 5 340 345 350
 Phe Arg Ala Thr Glu Ala Ala Lys Lys Ile His Leu Ser Asn Leu Ser
 355 360 365
 Arg Ile Asp Gln Glu Thr Gln Lys Ile Phe Lys Ala Gly Lys Leu Met
 10 370 375 380
 Lys Pro Asp Glu Leu Asp Ala Gln Ile Arg Asp Arg Ser Ala Arg Tyr
 385 390 395 400
 Val Gln Phe Asp Cys Phe His Glu Ile Leu Pro Gln Pro Glu Asn Arg
 15 405 410 415
 Ile Val Pro Ser Lys Thr Ala Thr Asp Ala Ile Gly Ile Pro Arg Pro
 420 425 430
 Glu Ile Thr Tyr Ala Ile Asp Asp Tyr Val Lys Arg Gly Ala Ala His
 20 435 440 445
 Thr Arg Glu Val Tyr Ala Thr Ala Ala Lys Val Leu Gly Gly Thr Asp
 450 455 460
 Val Val Phe Asn Asp Glu Phe Ala Pro Asn Asn His Ile Thr Gly Ser
 25 465 470 475 480
 Thr Ile Met Gly Ala Asp Ala Arg Asp Ser Val Val Asp Lys Asp Cys
 485 490 495
 Arg Thr Phe Asp His Pro Asn Leu Phe Ile Ser Ser Ser Ala Thr Met
 30 500 505 510
 Pro Thr Val Gly Thr Val Asn Val Thr Leu Thr Ile Ala Ala Leu Ala
 515 520 525
 Leu Arg Met Ser Asp Thr Leu Lys Lys Glu Val
 35 530 535

<210> 4

<211> 27

40 <212> PRT

<213> Burkholderia cepacia

<400> 4

45 Val Arg Lys Ser Thr Leu Thr Phe Leu Ile Ala Gly Cys Leu Ala Leu
 1 5 10 15
 Pro Gly Phe Ala Arg Ala Ala Asp Ala Ala Asp
 20 25

50

<210> 5

<211> 28

<212> DNA

55 <213> Artificial Sequence

EP 1 498 484 A1

<220>
5 <223> Description of Artificial Sequence: primer

<400> 5
10 cccaagcttg ggccgatacc gatacgca 28

<210> 6
<211> 29
<212> DNA
15 <213> Artificial Sequence

<220>
20 <223> Description of Artificial Sequence: primer

<400> 6
gagaagcttt ccgcacggtc agacttcc 29

25 <210> 7
<211> 27
<212> DNA
30 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

35 <400> 7
catgccatgg cacacaacga caacact 27

<210> 8
40 <211> 27
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence: primer

<400> 8
50 cccaagcttg ggtcagactt ccttcttcag c 31

<210> 9
<211> 16
55 <212> PRT

EP 1 498 484 A1

<213> Burkholderia cepacia

5 <400> 9
Ala Asp Ala Ala Asp Pro Ala Leu Val Lys Arg Gly Glu Tyr Leu Ala
1 5 10 15

10 <210> 10
<211> 25
<212> PRT
<213> Artificial Sequence

15 <220>
<223> Description of Artificial Sequence:consensus

20 <220>
<221> UNSURE
<222> (6, 17, 18, 19, 22)
<223> Xaa=unknown

25 <400> 10
Ala Asp Ala Ala Asp Xaa Ala Leu Val Lys Arg Gly Glu Tyr Leu Ala
1 5 10 15
30 Xaa Xaa Xaa Asp Cys Xaa Ala Cys His
20 25

35 <210> 11
<211> 2410
<212> DNA
<213> Burkholderia cepacia

40 <220>
<221> CDS
<222> (673)..(1950)

45 <400> 11
gatggaccac cggggcaccg gcggtgctgt ctacgcgaac gagaagctgt ggccggggccg 60
cgcccgccag gatacgacgt cgtgacggtt ttccgcgcac ggcccggttc gcgcgaccga 120
agcccggaag aagatccatc tgcgaacat gtcccgcatc aaccaggaga cgcagaagat 180
50 cltcaaggcc ggcaaactga tgaagcacga ggagctcgac gcgcagatcc gcgaccgttc 240
cgcgcgctac gtgcagttcg actgcttcca cgagatctcg ccgcagcccg agaaccgcat 300
cgigccgagc aagacggcca ccgacgcgat cgggaccccg cgcccgaga lcacgtatgc 360
gacgcagcat tacgtgaagc gcggcgccgt gcacacgcgc gaggctacg cgacggccgc 420
55 gaaggctcgt ggccggcaccg acgtcgtctt caacgacgag ttgcgcgcga acaaccacat 480

EP 1 498 484 A1

cacggcgcg aggaicalgg gcgcggatgc acgcgacicg gicgicgaca aggactgccg 540
 cacgticgac catccgaacc igticcicic gagcagcicg acgaigccga ccgtcggtac 600
 5 gglgaacglg acgclgacga icgcggcgct cgcgcicgcg atgicggaca cgtgaagaa 660
 ggaagtctga cc glg cgg aaa tct act ctc acc ttc ctc ctc gcc ggc tgc 711
 Val Arg Lys Ser Thr Leu Thr Phe Leu Leu Ala Gly Cys
 1 5 10
 10 ctc gcg ctg ccc ggc ctc gca cgc gcg gcc gat tgc gcc gat ccg gcg 759
 Leu Ala Leu Pro Gly Leu Ala Arg Ala Ala Asp Ser Ala Asp Pro Ala
 15 20 25
 cal gtc aag cgc ggc gaa tac ctc gcc gtc gcg ggc gac tgc atg gca 807
 15 His Val Lys Arg Gly Glu Tyr Leu Ala Val Ala Gly Asp Cys Met Ala
 30 35 40 45
 tgc cac acc gcg aag ggc ggc aag ccg ttc gcg ggc ggc ctc ggc atg 855
 Cys His Thr Ala Lys Gly Gly Lys Pro Phe Ala Gly Gly Leu Gly Met
 20 50 55 60
 ccg glg ccg atg ctc ggc aag atc taf acg agc aac atc aca ccg gat 903
 Pro Val Pro Met Leu Gly Lys Ile Tyr Thr Ser Asn Ile Thr Pro Asp
 65 70 75
 25 ccc gat acc ggc atc ggc aac igg acg ttc gag gac ttc gag cgc gcg 951
 Pro Asp Thr Gly Ile Gly Asn Trp Thr Phe Glu Asp Phe Glu Arg Ala
 80 85 90
 gtg cgg cac ggc gla tgc aag aac ggc gac aac ctg tac ccg gcg atg 999
 30 Val Arg His Gly Val Ser Lys Asn Gly Asp Asn Leu Tyr Pro Ala Met
 95 100 105
 ccg tac gtg tgc tac gcg aag atc aac gac gac gac gtg caa gcg ctg 1047
 Pro Tyr Val Ser Tyr Ala Lys Ile Asn Asp Asp Asp Val Gln Ala Leu
 35 110 115 120 125
 tac gcg tac ttc atg cac ggc gtc gaa ccg gtc aag cag gcg ccg ccg 1095
 Tyr Ala Tyr Phe Met His Gly Val Glu Pro Val Lys Gln Ala Pro Pro
 130 135 140
 40 aag aac gag atc ccc gcg ctg ctg agc atg cgc tgg ccg ctg aag atc 1143
 Lys Asn Glu Ile Pro Ala Leu Leu Ser Met Arg Trp Pro Leu Lys Ile
 145 150 155
 tgg aac tgg ctg ttc ctg aag gac ggc gtg tac cag ccg aag ccc gag 1191
 45 Trp Asn Trp Leu Phe Leu Lys Asp Gly Val Tyr Gln Pro Lys Pro Glu
 160 165 170
 cag agc gcc gag tgg aac cgc ggc gcc taf ctc gtg cag ggc ctc gcg 1239
 Gln Ser Ala Glu Trp Asn Arg Gly Ala Tyr Leu Val Gln Gly Leu Ala
 50 175 180 185
 cac tgc agc acg tgc cac acg ccg cgc ggc atc gcg atg cag gag aag 1287
 His Cys Ser Thr Cys His Thr Pro Arg Gly Ile Ala Met Gln Glu Lys
 190 195 200 205
 55 tcr ctc gac gaa acg ggc ggc agc ttc ctg tgc ggc tgc glg ctc gcg 1335

EP 1 498 484 A1

Ser Leu Asp Glu Thr Gly Gly Ser Phe Leu Ser Gly Ser Val Leu Ala
 210 215 220
 5 ggc tgg gac ggc tac aac atc acg tcc gac ccg aac ggc ggg atc ggc 1383
 Gly Trp Asp Gly Tyr Asn Ile Thr Ser Asp Pro Asn Ala Gly Ile Gly
 225 230 235
 10 ggc tgg acg cag cag cag ctc gtc cag tac ctg cgc acc ggc agc gtc 1431
 Gly Trp Thr Gln Gln Gln Leu Val Gln Tyr Leu Arg Thr Gly Ser Val
 240 245 250
 ccg ggc ctc ggc cag ggc gcc ggc ccg atg gcc gag ggc atc gag cac 1479
 Pro Gly Leu Ala Gln Ala Ala Gly Pro Met Ala Glu Ala Ile Glu His
 255 260 265
 15 agc ttc tgc aag atg acc gaa gcc gac atc ggc ggc ccg atg gcc gag 1527
 Ser Phe Ser Lys Met Thr Glu Ala Asp Ile Gly Gly Pro Met Ala Glu
 270 275 280 285
 20 ggc atc gag cac agc ttc tgc aag atg acc gaa gcc gac atc ggc cgc 1575
 Ala Ile Glu His Ser Phe Ser Lys Met Thr Glu Ala Asp Ile Gly Arg
 290 295 300
 tgc tgc tgg ggc aag ccg gcc gag gat ggc ctg aag ctg cgc ggc gtc 1623
 25 Ser Ser Trp Gly Lys Pro Ala Glu Asp Gly Leu Lys Leu Arg Gly Val
 305 310 315
 ggc ctc ggc tgc tgc ggc atc gat ccg gca ccg ctg tat ctc ggc aac 1671
 Ala Leu Ala Ser Ser Gly Ile Asp Pro Ala Pro Leu Tyr Leu Gly Asn
 320 325 330
 30 tgc ggc acc tgc cac cag atg cag ggc aag ggc acg ccg gac ggt tac 1719
 Cys Ala Thr Cys His Gln Met Gln Gly Lys Gly Thr Pro Asp Gly Tyr
 335 340 345
 35 tac ccg ccg ttg ttc cac aac tgc acg gtc ggc ggc tgc aat ccg acc 1767
 Tyr Pro Pro Leu Phe His Asn Ser Thr Val Gly Ala Ser Asn Pro Thr
 350 355 360 365
 40 aac ctc gtc cag gtc atc ctg aac ggc gtc cag cgc aag gcc ggc agc 1815
 Asn Leu Val Gln Val Ile Leu Asn Gly Val Gln Arg Lys Ala Gly Ser
 370 375 380
 gag gac gtc ggc atg ccc ggc ttc cgc cac gag ctg tgc gat ggc cag 1863
 Glu Asp Val Gly Met Pro Ala Phe Arg His Glu Leu Ser Asp Ala Gln
 385 390 395
 45 atc gcc ggc ctg acg aac tac ctg acg ggc cag ttc ggc aat ccg gcc 1911
 Ile Ala Ala Leu Thr Asn Tyr Leu Thr Gly Gln Phe Gly Asn Pro Ala
 400 405 410
 50 ggc aag gtc acc gag cag gac gtc ggc aag ctg cgc tga aacgcggcac 1960
 Ala Lys Val Thr Glu Gln Asp Val Ala Lys Leu Arg
 415 420 425
 55 gcggcgaggc agggcaacaa tagaaaagag gaggagcaca gcacatcggg caggccccga 2020
 tgcgggttgt tgcagagcgg gacggcgggc gcaggcggtc gcccgttcgt gttcataggc 2080

EP 1 498 484 A1

aatccggigc gcgcacgcgc cgcacgilit tegtigatcg agaccatgac accgaaccaa 2140
 ccgilitcgc cgtccagcgc cgaiglgcig ctcgicgtgt cccgaatccf gctcgtgac 2200
 5 cgtitcgtga iglicggcig gaagaagail atcgactict cccgtacga! cgcgticagt 2260
 ggcagcgagg gcgcgcgcgc gccgaicac tccggcgga tctccgtcgt gatggagctc 2320
 atcgtcggga ttcgatctt cgtcggtttc cagacgcgc cgcicgcgt gilgcttgcg 2380
 cgtiacacga tccgtaccgc calcatcggc 2410

10

<210> 12

<211> 425

<212> PRT

15

<213> Burkholderia cepacia

<400> 12

Val Arg Lys Ser Thr Leu Thr Phe Leu Leu Ala Gly Cys Leu Ala Leu
 1 5 10 15
 Pro Gly Leu Ala Arg Ala Ala Asp Ser Ala Asp Pro Ala His Val Lys
 20 25 30
 Arg Gly Glu Tyr Leu Ala Val Ala Gly Asp Cys Met Ala Cys His Thr
 25 35 40 45
 Ala Lys Gly Gly Lys Pro Phe Ala Gly Gly Leu Gly Met Pro Val Pro
 50 55 60
 Met Leu Gly Lys Ile Tyr Thr Ser Asn Ile Thr Pro Asp Pro Asp Thr
 30 65 70 75 80
 Gly Ile Gly Asn Trp Thr Phe Glu Asp Phe Glu Arg Ala Val Arg His
 85 90 95
 Gly Val Ser Lys Asn Gly Asp Asn Leu Tyr Pro Ala Met Pro Tyr Val
 35 100 105 110
 Ser Tyr Ala Lys Ile Asn Asp Asp Asp Val Gln Ala Leu Tyr Ala Tyr
 115 120 125
 Phe Met His Gly Val Glu Pro Val Lys Gln Ala Pro Pro Lys Asn Glu
 40 130 135 140
 Ile Pro Ala Leu Leu Ser Met Arg Trp Pro Leu Lys Ile Trp Asn Trp
 145 150 155 160
 Leu Phe Leu Lys Asp Gly Val Tyr Gln Pro Lys Pro Glu Gln Ser Ala
 45 165 170 175
 Glu Trp Asn Arg Gly Ala Tyr Leu Val Gln Gly Leu Ala His Cys Ser
 180 185 190
 Thr Cys His Thr Pro Arg Gly Ile Ala Met Gln Glu Lys Ser Leu Asp
 50 195 200 205
 Glu Thr Gly Gly Ser Phe Leu Ser Gly Ser Val Leu Ala Gly Trp Asp
 210 215 220
 Gly Tyr Asn Ile Thr Ser Asp Pro Asn Ala Gly Ile Gly Gly Trp Thr
 55 225 230 235 240

EP 1 498 484 A1

Gln Gln Gln Leu Val Gln Tyr Leu Arg Thr Gly Ser Val Pro Gly Leu
 245 250 255
 5 Ala Gln Ala Ala Gly Pro Met Ala Glu Ala Ile Glu His Ser Phe Ser
 260 265 270
 Lys Met Thr Glu Ala Asp Ile Gly Gly Pro Met Ala Glu Ala Ile Glu
 275 280 285
 10 His Ser Phe Ser Lys Met Thr Glu Ala Asp Ile Gly Arg Ser Ser Trp
 290 295 300
 Gly Lys Pro Ala Glu Asp Gly Leu Lys Leu Arg Gly Val Ala Leu Ala
 305 310 315 320
 15 Ser Ser Gly Ile Asp Pro Ala Pro Leu Tyr Leu Gly Asn Cys Ala Thr
 325 330 335
 Cys His Gln Met Gln Gly Lys Gly Thr Pro Asp Gly Tyr Tyr Pro Pro
 340 345 350
 20 Leu Phe His Asn Ser Thr Val Gly Ala Ser Asn Pro Thr Asn Leu Val
 355 360 365
 Gln Val Ile Leu Asn Gly Val Gln Arg Lys Ala Gly Ser Glu Asp Val
 370 375 380
 25 Gly Met Pro Ala Phe Arg His Glu Leu Ser Asp Ala Gln Ile Ala Ala
 385 390 395 400
 Leu Thr Asn Tyr Leu Thr Gly Gln Phe Gly Asn Pro Ala Ala Lys Val
 405 410 415
 30 Thr Glu Gln Asp Val Ala Lys Leu Arg
 420 425

35 <210> 13
 <211> 27
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence: primer

45 <400> 13
 tgcaccgigc ggaaatciac tctcact

27

50 <210> 14
 <211> 27
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> Description of Artificial Sequence: primer

EP 1 498 484 A1

<400> 14
 5 actlcccllcl lcagcgtgic cgacatc 27

 <210> 15
 <211> 1441
 10 <212> DNA
 <213> Burkholderia cepacia

 <220>
 15 <221> CDS
 <222> (121).. (1398)

 <400> 15
 20 iccgaaccig ttcattlccga gcagcgcgac gattcccgacc gtcggtaccg taaacgigac 60
 gcigacgalt gccgcgcicg cgcicggai gtcggacacg ctgaagaagg aagtcigacc 120
 glg cgg aaa tct act ctc act ttc ctc atc gcc ggc tgc ctc gcg ilg 168
 Val Arg Lys Ser Thr Leu Thr Phe Leu Ile Ala Gly Cys Leu Ala Leu
 25 1 5 10 15
 ccg ggc ttc gcg cgc gcg gcc gat gcg gcc gat ccg gcg ctg gtc aag 216
 Pro Gly Phe Ala Arg Ala Ala Asp Ala Ala Asp Pro Ala Leu Val Lys
 20 25 30
 30 cgc ggc gaa tac ctc gcg acc gcc atg ccg gta ccg atg ctc ggc aag 264
 Arg Gly Glu Tyr Leu Ala Thr Ala Met Pro Val Pro Met Leu Gly Lys
 35 35 40 45
 atc tac acg agc aac atc acg ccc gat ccc gat acg ggc gac tgc atg 312
 Ile Tyr Thr Ser Asn Ile Thr Pro Asp Pro Asp Thr Gly Asp Cys Met
 50 55 60
 gcc tgc cac acc glg aag ggc ggc aag ccg tac gcg ggc ggc ctt ggc 360
 Ala Cys His Thr Val Lys Gly Gly Lys Pro Tyr Ala Gly Gly Leu Gly
 40 65 70 75 80
 ggc atc ggc aaa tgg acg ttc gag gac ttc gag cgc gcg gtg cgg cac 408
 Gly Ile Gly Lys Trp Thr Phe Glu Asp Phe Glu Arg Ala Val Arg His
 85 90 95
 45 ggc glg tgc aag aac ggc gac aac ctg tat ccg gcg atg ccg tac glg 456
 Gly Val Ser Lys Asn Gly Asp Asn Leu Tyr Pro Ala Met Pro Tyr Val
 100 105 110
 50 tgc tac gcg aag atc aag gac gac gac gla cgc gcg ctg tac gcc tac 504
 Ser Tyr Ala Lys Ile Lys Asp Asp Val Arg Ala Leu Tyr Ala Tyr
 115 120 125
 ttc atg cac ggc glc gag ccg gtc aag cag gcg ccg ccg aag aac gag 552
 55 Phe Met His Gly Val Glu Pro Val Lys Gln Ala Pro Pro Lys Asn Glu
 130 135 140

EP 1 498 484 A1

aic cca gcg ctg cta agc atg cgc tgg ccg ctg aag atc tgg aac tgg 600
 Ile Pro Ala Leu Leu Ser Met Arg Trp Pro Leu Lys Ile Trp Asn Trp
 145 150 155 160
 .5 ctg ttc ctg aag gac ggc ccg tac cag ccg aag ccg tgg cag agc gcc 648
 Leu Phe Leu Lys Asp Gly Pro Tyr Gln Pro Lys Pro Ser Gln Ser Ala
 165 170 175
 10 gaa tgg aat cgc ggc gcg tat ctg gig cag ggt ctc gcg cac tgc agc 696
 Glu Trp Asn Arg Gly Ala Tyr Leu Val Gln Gly Leu Ala His Cys Ser
 180 185 190
 acg tgc cac acg ccg cgc ggc atc gcg atg cag gag aag tgc ctc gac 744
 15 Thr Cys His Thr Pro Arg Gly Ile Ala Met Gln Glu Lys Ser Leu Asp
 195 200 205
 gaa acc ggc ggc agc ttc ctc gcg ggg tgg gig ctc gcc ggc tgg gac 792
 Glu Thr Gly Gly Ser Phe Leu Ala Gly Ser Val Leu Ala Gly Trp Asp
 210 215 220
 20 ggc tac aac atc acg tgc gac ccg aat gcg ggg atc ggc agc tgg acc 840
 Gly Tyr Asn Ile Thr Ser Asp Pro Asn Ala Gly Ile Gly Ser Trp Thr
 225 230 235 240
 25 cag cag cag ctc gig cag tat tlg cgc acc ggc agc gig ccg ggc gic 888
 Gln Gln Gln Leu Val Gln Tyr Leu Arg Thr Gly Ser Val Pro Gly Val
 245 250 255
 gcg cag gcg gcc ggg ccg atg gcc gag gcg gtc gag cac agc ttc tgc 936
 30 Ala Gln Ala Ala Gly Pro Met Ala Glu Ala Val Glu His Ser Phe Ser
 260 265 270
 aag atg acc gaa gcg gac atc ggt gcg atc gcc acc tac gtc cgc acc 984
 Lys Met Thr Glu Ala Asp Ile Gly Ala Ile Ala Thr Tyr Val Arg Thr
 275 280 285
 35 gig ccg gcc gtt gcc gac agc aac gcg aag cag ccg ccg tgc tgc tgg 1032
 Val Pro Ala Val Ala Asp Ser Asn Ala Lys Gln Pro Arg Ser Ser Trp
 290 295 300
 40 ggc aag ccg gcc gag gac ggg ctg aag ctg cgc ggt gtc gcg ctc gcg 1080
 Gly Lys Pro Ala Glu Asp Gly Leu Lys Leu Arg Gly Val Ala Leu Ala
 305 310 315 320
 tgc tgc ggc atc gat ccg gcg ccg ctg tat ctc ggc aac tgc gcg acc 1128
 45 Ser Ser Gly Ile Asp Pro Ala Arg Leu Tyr Leu Gly Asn Cys Ala Thr
 325 330 335
 tgc cac cag atg cag ggc aag ggc acc ccg gac ggc tat tac ccg tgc 1176
 Cys His Gln Met Gln Gly Lys Gly Thr Pro Asp Gly Tyr Tyr Pro Ser
 340 345 350
 50 ctg ttc cac aac tcc acc gtc ggc gcg tgc aat ccg tgc aac ctc gtg 1224
 Leu Phe His Asn Ser Thr Val Gly Ala Ser Asn Pro Ser Asn Leu Val
 355 360 365
 55 cag gig atc ctg aac ggc gig cag cgc aag atc ggc agc gag gat atc 1272

EP 1 498 484 A1

Gln Val Ile Leu Asn Gly Val Gln Arg Lys Ile Gly Ser Glu Asp Ile
 370 375 380
 5 ggg atg ccc gct ttc cgc tac gat ctc aac gac gcg cag atc gcc gcg 1320
 Gly Met Pro Ala Phe Arg Tyr Asp Leu Asn Asp Ala Gln Ile Ala Ala
 385 390 395 400
 10 ctg acg aac tac gtc acc gcg cag ttc ggc aat ccg gcg gcg aag gtc 1368
 Leu Thr Asn Tyr Val Thr Ala Gln Phe Gly Asn Pro Ala Ala Lys Val
 405 410 415
 acg gag cag gac gtc gcg aag ctg cgc tga catagtcggg cgcgcgcgaca 1418
 Thr Glu Gln Asp Val Ala Lys Leu Arg
 15 420 425
 cggcgcaacc galaggacag gag 1441
 <210> 16
 20 <211> 425
 <212> PRT
 <213> Burkholderia cepacia
 25 <400> 16
 Val Arg Lys Ser Thr Leu Thr Phe Leu Ile Ala Gly Cys Leu Ala Leu
 5 10 15
 30 Pro Gly Phe Ala Arg Ala Ala Asp Ala Ala Asp Pro Ala Leu Val Lys
 20 25 30
 Arg Gly Glu Tyr Leu Ala Thr Ala Met Pro Val Pro Met Leu Gly Lys
 35 40 45
 35 Ile Tyr Thr Ser Asn Ile Thr Pro Asp Pro Asp Thr Gly Asp Cys Met
 50 55 60
 Ala Cys His Thr Val Lys Gly Gly Lys Pro Tyr Ala Gly Gly Leu Gly
 65 70 75 80
 40 Gly Ile Gly Lys Trp Thr Phe Glu Asp Phe Glu Arg Ala Val Arg His
 85 90 95
 Gly Val Ser Lys Asn Gly Asp Asn Leu Tyr Pro Ala Met Pro Tyr Val
 100 105 110
 45 Ser Tyr Ala Lys Ile Lys Asp Asp Asp Val Arg Ala Leu Tyr Ala Tyr
 115 120 125
 Phe Met His Gly Val Glu Pro Val Lys Gln Ala Pro Pro Lys Asn Glu
 130 135 140
 50 Ile Pro Ala Leu Leu Ser Met Arg Trp Pro Leu Lys Ile Trp Asn Trp
 145 150 155 160
 Leu Phe Leu Lys Asp Gly Pro Tyr Gln Pro Lys Pro Ser Gln Ser Ala
 165 170 175
 55 Glu Trp Asn Arg Gly Ala Tyr Leu Val Gln Gly Leu Ala His Cys Ser
 180 185 190

EP 1 498 484 A1

Thr Cys His Thr Pro Arg Gly Ile Ala Met Gln Glu Lys Ser Leu Asp
 195 200 205
 5 Glu Thr Gly Gly Ser Phe Leu Ala Gly Ser Val Leu Ala Gly Trp Asp
 210 215 220
 Gly Tyr Asn Ile Thr Ser Asp Pro Asn Ala Gly Ile Gly Ser Trp Thr
 225 230 235 240
 10 Gln Gln Gln Leu Val Gln Tyr Leu Arg Thr Gly Ser Val Pro Gly Val
 245 250 255
 Ala Gln Ala Ala Gly Pro Met Ala Glu Ala Val Glu His Ser Phe Ser
 260 265 270
 15 Lys Met Thr Glu Ala Asp Ile Gly Ala Ile Ala Thr Tyr Val Arg Thr
 275 280 285
 Val Pro Ala Val Ala Asp Ser Asn Ala Lys Gln Pro Arg Ser Ser Trp
 290 295 300
 20 Gly Lys Pro Ala Glu Asp Gly Leu Lys Leu Arg Gly Val Ala Leu Ala
 305 310 315 320
 Ser Ser Gly Ile Asp Pro Ala Arg Leu Tyr Leu Gly Asn Cys Ala Thr
 325 330 335
 25 Cys His Gln Met Gln Gly Lys Gly Thr Pro Asp Gly Tyr Tyr Pro Ser
 340 345 350
 Leu Phe His Asn Ser Thr Val Gly Ala Ser Asn Pro Ser Asn Leu Val
 355 360 365
 30 Gln Val Ile Leu Asn Gly Val Gln Arg Lys Ile Gly Ser Glu Asp Ile
 370 375 380
 Gly Met Pro Ala Phe Arg Tyr Asp Leu Asn Asp Ala Gln Ile Ala Ala
 385 390 395 400
 35 Leu Thr Asn Tyr Val Thr Ala Gln Phe Gly Asn Pro Ala Ala Lys Val
 405 410 415
 Thr Glu Gln Asp Val Ala Lys Leu Arg
 420 425
 40
 <210> 17
 <211> 5
 <212> PRT
 45 <213> Artificial Sequence
 <220>
 50 <223> Description of Artificial Sequence: heme binding motif
 <220>
 <221> UNSURE
 <222> (2, 3)
 55 <223> Xaa=unknown

EP 1 498 484 A1

<400> 17

Cys Xaa Xaa Cys His

1 5

<210> 18

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 18

caigccalgg cacacaacga caacact

27

<210> 19

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 19

cccaagcttg ggtcagactt ccttcctcag c

31

Claims

1. A protein defined in the following (A) or (B):

(A) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16;

(B) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16 including substitution, deletion, insertion or addition of 1 to 20 amino acid residues.

50

2. A DNA encoding a protein defined in the following (A) or (B):

(A) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16;

(B) a protein which has at least the amino acid sequence comprising amino acids 23 to 425 of SEQ ID NO: 16 including substitution, deletion, insertion or addition of 1 to 20 amino acid residues.

55

3. The DNA according to claim 2, wherein the DNA is defined in the following (a) or (b):

EP 1 498 484 A1

(a) a DNA comprising the nucleotide sequence consisting of nucleotides 187 to 1398 of SEQ ID NO: 15;
(b) a DNA which is hybridizable with the nucleotide sequence consisting of nucleotides 187 to 1398 of SEQ ID NO: 15 under stringent conditions.

- 5 4. The DNA according to claim 3, further comprising the nucleotide sequence consisting of nucleotides 121 to 187 of SEQ ID NO: 15.
5. A recombinant vector comprising the DNA according to any one of claims 2 to 4.
- 10 6. A transformant transformed with the DNA according to any one of claims 2 to 4 or the recombinant vector according to claim 5.
7. A method of producing a glucose dehydrogenase β subunit, comprising culturing the transformant according to claim 6 to produce a glucose dehydrogenase β subunit as an expression product of the DNA, and collecting the produced β subunit.
- 15 8. The DNA according to claim 3 or 4, further comprising the nucleotide sequence encoding an α subunit and a γ subunit of glucose dehydrogenase of *Burkholderia cepacia*.
- 20 9. A recombinant vector comprising the DNA according to claim 8.
10. A transformant transformed with the DNA according to claim 8 or the recombinant vector according to claim 9.
- 25 11. A method of producing a glucose dehydrogenase complex, comprising culturing the transformant according to claim 10 to produce a glucose dehydrogenase complex as an expression product of the DNA, and collecting the produced complex.
- 30
- 35
- 40
- 45
- 50
- 55

EP 1 498 484 A1

INTERNATIONAL SEARCH REPORT

International application No.
ECT/JP03/05375

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C12N15/09, 9/04, 1/15, 1/19, 1/21, 5/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C12N15/09, 9/04, 1/15, 1/19, 1/21, 5/00		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) SwissProt/PIR/GeneSeq GenBank/EMBL/DBJ/GeneSeq BIOSIS/WPI (DIALOG)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SODE, K. et al., A novel thermostable glucose dehydrogenase varying temperature properties by altering its quaternary structures Enzyme Microb. Technol., 1996, 19(2), pages 82 to 85	1-11
P, X	WO 02/36779 A1 (Koji SODE), 10 May, 2002 (10.05.02), Full text (Family: none)	1-11
A	YAMAZAKI, T. et al., Subunit analyses of a novel thermostable glucose dehydrogenase showing different temperature properties according to its quaternary structure Appl. Biochem. Biotechnol., 1999, 77-79, pages 325 to 335	1-11
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 25 June, 2003 (25.06.03)		Date of mailing of the international search report 15 July, 2003 (15.07.03)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

EP 1 498 484 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/05375

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	YAMAZAKI, T. et al., Increased thermal stability of glucose dehydrogenase by cross-linking chemical modification <i>Biotechnol.lett.</i> , 1999, 21(3), pages 199 to 202	1-11
P,A	INOSE, K. et al., Cloning and expression of the gene encoding catalytic subunit of thermostable glucose dehydrogenase from <i>Burkholderia cepacia</i> in <i>Escherichia coli</i> <i>Biochim.Biophys.Acta.</i> , 2003 February, 1645(2), p.133-8	1-11

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.